

50 LBS. (22.68) NET WEIGHT

**NO. 10132****AMPROLIUM CRUMBLES****FOR BEEF AND DAIRY CALVES****MEDICATED**

Feed as an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed as an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

**ACTIVE DRUG INGREDIENT**

Amprolium . . . . . 2.27 grams/pound

**GUARANTEED ANALYSIS**

Crude Protein . . . . . Minimum . . . . . 14.0%

Crude Fat . . . . . Minimum . . . . . 3.5%

Crude Fiber . . . . . Maximum . . . . . 25.0%

**INGREDIENTS**

Processed grain by-products, grain products, molasses products, dried lignin sulfonate.

**FEEDING DIRECTIONS**

**Prevention:** Feed 0.1 lbs. per 100 lbs. body weight to furnish 227 mg/100 lbs. body weight. Continue feeding for 21 days.

**Treatment:** Feed 0.2 lbs. per 100 lbs. body weight to furnish 454 mg/100 lbs. body weight for 5 days. Treatment should be administered on a herd basis. When one calf shows coccidiosis, it is likely that the rest of the group has been exposed and all calves in the group should be treated.

**WARNING:** Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not use in Type B or Type C medicated feeds containing bentonite.

**LIMITATIONS:** Top-dress on or mix in the daily ration to provide 5 milligrams per kilogram of body weight per day. Feed for 21 days when experience indicates that coccidiosis is likely to be a hazard, as the sole source of Amprolium.



**Manufactured By**  
**Hi-Pro Feeds LLC**  
 Home Office  
 P.O. Box 519  
 Friona, Texas 79035