

**PRECAUTIONARY STATEMENTS:**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING:** This product contains high levels of copper. Consumption of this product by sheep or goats may result in copper toxicity. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID:** If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-764-7661 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS:** Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**FEEDING AND MANAGEMENT INSTRUCTIONS**

When used as directed, Cowman's Choice IGR Fly Control Mineral will provide sufficient S-Methoprene Insect Growth Regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of Cowman's Choice IGR Fly Control Mineral per 100 pounds of animal body weight per day, which is equivalent to 4 ounces per day for a 1000 pound animal.

1. Assure cattle are not starved for nutrients in Cowman's Choice IGR Fly Control Mineral by providing them prior to feeding Cowman's Choice IGR Fly Control Mineral.
2. Offer cattle 50 lbs. of Cowman's Choice IGR Fly Control Mineral for every 15 head of cattle.
3. Remove all other self limiting nutritional supplements.
4. Locate Cowman's Choice Fly Control Mineral where cattle congregate (watering, loafing, shade areas).
5. Replenish Cowman's Choice IGR Fly Control Mineral on a regular basis as needed.
6. If intake of Cowman's Choice IGR Fly Control Mineral is below 0.4 ounces per 100 pounds animal body weight per day, increase the number of feeding locations and/or change location(s) to areas more frequented by cattle.
7. If intake of Cowman's Choice IGR Fly Control Mineral is above 0.4 ounces per 100 pounds animal body weight per day, reduce the number of feeding locations and/or change location(s) to areas less frequented by cattle.
8. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For information, call 1-800-447-7620 or visit our website: [hiprofeeds.com](http://hiprofeeds.com)  
Altosid® Insect Growth Regulator is a registered trademark of Wellmark International.  
EPA REG. NO. 2724-494-26171  
EPA EST. NO. 26171-NM-1



**Manufactured By:**  
**Hi-Pro Feeds LLC**  
Home Office  
P.O. Box 519  
Frona, Texas 79035

**BACKSIDE #10513**

10-16-19