

10545

9/19

50 LBS. (22.68 KG) NET WEIGHT

NO. 10545

PRIMERO RED GOAT FEED

FOR GROWING GOATS

MEDICATED

For the prevention of Coccidiosis in young goats caused by *Eimeria christenseni* and *Eimeria ninakohlyakimovae*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.2 Grams Per Ton

GUARANTEED ANALYSIS

Crude Protein	Minimum	16.0%
(This includes not more than 1.2% equivalent non-protein nitrogen.)		
Crude Fat	Minimum	1.8%
Crude Fiber	Maximum	21.0%
Calcium	Minimum	1.0%
Calcium	Maximum	1.5%
Phosphorus	Minimum	0.3%
Salt	Minimum	1.0%
Salt	Maximum	1.5%
Copper	Minimum	18.0 ppm
Copper	Maximum	35.0 ppm
Selenium	Minimum	0.2 ppm
Vitamin A	Minimum	3,000 IU/LB

INGREDIENTS

Roughage products (40%), grain products, forage products, plant protein products, processed grain by-products, molasses products, calcium carbonate, salt, ammonium chloride, calcium sulfate, silicon dioxide, zinc sulfate, copper sulfate, Vitamin E supplement, manganese sulfate, cobalt sulfate, Vitamin A supplement, selenium, Vitamin D supplement, magnesium oxide, zinc amino acid complex, manganese amino acid complex, copper amino acid complex, cobalt glucoheptonate, natural and artificial flavoring and sodium selenite.

FEEDING DIRECTIONS

Feed at the rate of 1.67 pounds per 100 pounds body weight to provide 22.7 mg/100 lb. (0.5mg/kg) body weight of Decoquinatate per day. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Coccidiostats are not indicated for use in adult animals due to continuous previous exposure. Always provide plenty of clean fresh water for drinking.

CAUTION: Do not feed to goats producing milk for food.

WARNING: Do not use in feeds containing bentonite.



Manufactured By:
Hi-Pro Feeds LLC
 Home Office
 P.O. Box 519
 Friona, Texas 79035