

13395375

8/21

50 LBS. (22.68) NET WEIGHT

NO. 13395375

AMPROLIUM CRUMBLES

FOR BEEF AND DAIRY CALVES

MEDICATED

Feed as an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed as an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Amprolium 2.27 grams/pound

GUARANTEED ANALYSIS

Crude Protein Minimum 14.0%

Crude Fat Minimum 3.5%

Crude Fiber Maximum 25.0%

INGREDIENTS

Processed grain by-products, grain products, molasses products, dried lignin sulfonate.

FEEDING DIRECTIONS

Prevention: Feed 0.1 lbs. per 100 lbs. body weight to furnish 227 mg/100 lbs. body weight. Continue feeding for 21 days.

Treatment: Feed 0.2 lbs. per 100 lbs. body weight to furnish 454 mg/100 lbs. body weight for 5 days. Treatment should be administered on a herd basis. When one calf shows coccidiosis, it is likely that the rest of the group has been exposed and all calves in the group should be treated.

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use in Type B or Type C medicated feeds containing bentonite.

LIMITATIONS: Top-dress on or mix in the daily ration to provide 5 milligrams per kilogram of body weight per day. Feed for 21 days when experience indicates that coccidiosis is likely to be a hazard, as the sole source of Amprolium.



Manufactured By
Hi-Pro Feeds LLC
Home Office
P.O. Box 519
Frona, Texas 79035