

13395425

6/23

50 LBS. (22.68 KG) NET WEIGHT
IGR 2:1 FLY CONTROL MINERAL
FOR BEEF CATTLE ON PASTURE

Prevents adult horn fly emergence from manure of treated cattle. A nutritional supplement containing S-Methoprene* insect growth regulator for continuous feeding to cattle through the horn fly season.

ACTIVE INGREDIENT

S-Methoprene (CAS #65733-16-6)	0.01%
OTHER INGREDIENTS	99.99%
Total	100.00%

*As the insect growth regulator Altosid®

GUARANTEED ANALYSIS

Calcium	Minimum	13.8%
Calcium	Maximum	15.7%
Phosphorus	Minimum	7.0%
Salt	Minimum	23.1%
Salt	Maximum	26.2%
Magnesium	Minimum	2.0%
Potassium	Minimum	1.0%
Copper	Minimum	1,800.0 ppm
Selenium	Minimum	30.0 ppm
Zinc	Minimum	5,000.0 ppm
Vitamin A	Minimum	150,000 IU/LB

INGREDIENTS

Monocalcium phosphate, salt, calcium carbonate, processed grain by-products, magnesium oxide, cane molasses, vegetable oil, lecithin, phosphoric acid, sulfuric acid, natural and artificial flavors, preservatives, propionic acid and sodium benzoate, potassium chloride, sodium selenite, zinc sulfate, copper sulfate, manganese sulfate, ethylenediamine dihydriodide, ferrous carbonate, mineral oil, magnesium oxide and cobalt carbonate, iron oxide, xtra-dry, Vitamin A supplement, Vitamin D supplement, Vitamin E supplement.

KEEP OUT OF REACH OF CHILDREN

WARNING:

SEE PRECAUTIONARY STATEMENTS

SEE FEEDING DIRECTIONS ON BACK

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222-24 hrs/day, 7 days/wk for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: This product is to be used as a cattle feed-through pesticide for the control of horn flies. All feed ingredients listed on this product label must adhere to the specifications and limits (if applicable) ascribed by the Food and Drug Administration (FDA) for the use of pesticides in cattle feed. This product contains ethoxyquin, a preservative. The amount of ethoxyquin may vary according to the individual manufacturer's formulation and will not exceed the 150 ppm maximum limit established by ethoxyquin by FDA, per 21 CFR 573.380

FEEDING AND MANAGEMENT INSTRUCTIONS

Feed this product to cattle only. When used as directed, IGR 2:1 Fly Control Mineral will provide sufficient S-Methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of IGR 2:1 Fly Control Mineral per 100 pounds of cattle body weight per day, which is equivalent to 4 ounces per day for a 1000 pound cow.

1. Assure cattle are not starved for nutrients in IGR 2:1 Fly Control Mineral by providing them prior to feeding IGR 2:1 Fly Control Mineral.
2. This product contains added copper. Do not feed to cattle in close proximity to sheep or other species that have a low tolerance to supplemental copper.
3. Offer cattle 50 lbs. of IGR2:1 Fly Control Mineral for every 15 head of cattle.
4. Remove all other self-limiting nutritional supplements while cattle are consuming this product.
5. Locate IGR 2:1 Fly Control Mineral where cattle congregate (watering, loafing, shade areas).
6. Replenish IGR 2:1 Fly Control Mineral on a regular basis as needed.
7. If intake of IGR 2:1 Fly Control Mineral is below 0.4 ounces per 100 pounds cattle body weight per day, reduce the number of feeding locations and/or change location(s) to areas more frequented by cattle.
8. If intake of IGR 2:1 Fly Control Mineral is above 0.4 ounces per 100 pounds animal body weight per day, reduce the number of feeding locations and/or change location(s) to areas less frequented by cattle.
9. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if eliminations of the adult fly population is desired.

For information call 1-800-447-7620.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then offer for recycling, if available, or, dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



Manufactured By:
Hi-Pro Feeds LLC
Home Office
P.O. Box 519
Friona, Texas 79035