

NO. 13399435

DAIRY CALF STARTER 19%

FOR YOUNG DAIRY CALVES.

MEDICATED

For the prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii* in calves (excluding veal calves).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 60.5 g/ton

Guaranteed Analysis

Crude Protein	(Min.)	19.0	%
Crude Fat	(Min.)	2.5	%
Crude Fiber	(Max.)	5.0	%
ADF	(Max.)	6.3	%
Calcium	(Min.)	0.5	%
Calcium	(Max.)	1.0	%
Phosphorus	(Min.)	0.4	%
Selenium	(Min.)	0.3	ppm
Vitamin A	(Min.)	12,000	IU/LB

Ingredients

GRAIN PRODUCTS, PLANT PROTEIN PRODUCTS, MOLASSES PRODUCTS, GLYCERIN, VEGETABLE OIL REFINERY LIPID WITH LECITHIN, PROCESSED GRAIN BY-PRODUCTS, CALCIUM CARBONATE, SALT, CALCIUM SULFATE, ROUGHAGE PRODUCTS, VITAMIN A SUPPLEMENT, MINERAL OIL, VITAMIN E SUPPLEMENT, ZINC PROTEINATE, ZINC SULFATE, COPPER SULFATE, MANGANESE SULFATE, ETHYLENEDIAMINE DIHYDRIODIDE, FERROUS CARBONATE, SODIUM SELENITE, MAGNESIUM OXIDE, COBALT CARBONATE, SODIUM SACCHARIN, ETHYL BURYRATE, ETHANOL, ETHYL PROPIONATE, CHOLINE CHLORIDE, VITAMIN D3 SUPPLEMENT, RIBOFLAVIN, D-CALCIUM PANTOTHENATE, NIACIN SUPPLEMENT, VITAMIN B12 SUPPLEMENT, THIAMINE MONONITRATE, BIOTIN, PYRIDOXINE HYDROCHLORIDE, MENADIONE SODIUM BISULFITE COMPLEX (SOURCE OF VITAMIN K ACTIVITY), FOLIC ACID, COPPER PROTEINATE AND VITAMIN D SUPPLEMENT.

Feeding Directions

Feed to calves at a rate to provide 0.14-1.0 mg/lb. of bodyweight of Monensin per day and up to 200 mg/head/day depending on the severity of the challenge. Consult your nutritionist for specific feeding directions. Always provide plenty of clean fresh water for drinking.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal.

Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

CAUTION: THIS FEED CONTAINS ADDED COPPER. DO NOT FEED TO SHEEP OR OTHER RELATED SPECIES.

Manufactured By:
Hi-Pro Feeds LLC
Home Office
P.O. Box 519
Friona, Texas 79035
Bulk
13399435 07/25

