

NO. 13402015

16% SHEEP & GOAT PELLET

FOR GROWING GOATS.

MEDICATED

For the prevention of coccidiosis in young sheep caused by Eimeria ovinoidalis, E. crandallis, E. parva and E. bakuensis.

ACTIVE DRUG INGREDIENT

Decoquinatate 27 g/ton

Guaranteed Analysis

Crude Protein*	(Min.)	16.0	%
* (This includes not more than 0.8 % equivalent non-protein nitrogen.)			
Crude Fat	(Min.)	3.0	%
Crude Fiber	(Max.)	16.5	%
Acid Detergent Fiber	(Max.)	23.8	%
Calcium	(Min.)	1.0	%
Calcium	(Max.)	1.5	%
Salt	(Min.)	0.5	%
Salt	(Max.)	1.0	%
Copper	(Min.)	9.0	ppm
Copper	(Max.)	10.0	ppm
Vitamin A	(Min.)	9,000	IU/LB

Ingredients

PROCESSED GRAIN BY-PRODUCTS, ROUGHAGE PRODUCTS, PLANT PROTEIN PRODUCTS, CALCIUM CARBONATE, MOLASSES PRODUCTS, DRIED LIGNIN SULFONATE, SALT, AMMONIUM CHLORIDE, ZINC SULFATE, MANGANESE SULFATE, MAGNESIUM OXIDE, ETHYLENEDIAMINE DIHYDRIDE, COBALT SULFATE, SODIUM SELENITE, FERROUS CARBONATE, IRON OXIDE, VITAMIN A SUPPLEMENT, VITAMIN D SUPPLEMENT AND VITAMIN E SUPPLEMENT.

Feeding Directions

Feed to goats as a complete feed at the rate of 3.0% to 4.0% of their body weight for growth. Always provide plenty of clean fresh water for drinking.

WARNING: Do not feed to sheep producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Feed Type C feed or milk replacer at a rate to provide 22.7 mg decoquinatate per 100 lb (0.5 mg/kg) bodyweight. Type C feed may be prepared from dry or liquid type B feed.

CAUTION: USE ONLY AS DIRECTED. FOR RUMINANTS (CATTLE, SHEEP AND GOATS) ONLY.

Manufactured By:

Hi-Pro Feeds LLC

Home Office

P.O. Box 519

Friona, Texas 79035

50 lbs (22.68 KG) NET WEIGHT

13402015 06/21

