



INSTANT DOE'S MATCH® KID MILK REPLACER

GUARANTEED ANALYSIS

Crude Protein, not less than	25.0%
Crude Fat, not less than	28.0%
Crude Fiber, not more than	0.15%
Calcium (Ca), not less than	0.75%
Calcium (Ca), not more than	1.25%
Phosphorus (P), not less than	0.70%
Sodium (Na), not more than	1.3%
Selenium (Se), not less than	0.3 ppm
Vitamin A, not less than	20,000 IU/lb
Vitamin D ₃ , not less than	5,000 IU/lb
Vitamin E, not less than	150 IU/lb

INGREDIENTS

Dried Whey Protein Concentrate, Dried Whey, Dried Whey Product, Dried Skimmed Milk, Dried Milk Protein, Animal Fat (preserved with BHA and BHT), Lecithin, Yeast Extract, Brewers Dried Yeast, Hydrolyzed Yeast, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Thiamine Mononitrate, Ascorbic Acid, Calcium Pantothenate, Niacin Supplement, Riboflavin, Biotin, Menadione Sodium Bisulfite Complex (source of Vitamin K₃), Pyridoxine Hydrochloride, Folic Acid, Vitamin B₁₂ Supplement, Choline Chloride, Fructooligosaccharides, Calcium Silicate, Potassium Sorbate (a preservative), Malic Acid, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Ethylenediamine Dihydroiodide and Selenium Yeast.

Doe's Match and LAND O LAKES logo are registered trademarks of Land O'Lakes, Inc.

To be fed to goat kids from second day of life through weaning.

MIXING DIRECTIONS

Always weigh Doe's Match Kid Milk Replacer powder for accurate mixing. Use **ONLY LOW SODIUM (<50 ppm)** water for mixing and feeding!

While stirring, mix 0.37 lbs dry weight or fill enclosed measure* heaping full of Doe's Match Kid Milk Replacer added to 2 pints of 110-120° F. water by slowly adding powder to water. Mix thoroughly.

*Enclosed measure holds approximately 0.37 lbs dry weight when heaping full. Weigh dry milk replacer powder for accurate mixing.

HAND FEEDING GUIDELINES

NOTE: ALWAYS offer free choice **LOW SODIUM (<50 ppm)** water.

Use the following schedule as a guideline for feeding reconstituted milk replacer:

Kid Weight (lbs)	Milk Replacer Powder Fed Per Day (lbs)	Pints of Reconstituted Milk Replacer Fed Per Day*
6	0.2	1.20
9	0.3	1.80
12	0.4	2.40
15	0.5	3.00

*Daily amounts should be split into 3 to 4 feedings daily at first, feeding no more than 1/2 pint per feeding. Feeding rates may be adjusted according to age and desired growth goals. **Always offer free choice water and high quality dry feed.**

FREE CHOICE FEEDING

Use of a free choice device requires that the milk replacer solution be kept cool (< 60° F.) to prevent overconsumption. Kids may need to be trained to suckle from nipples; however, they will readily adapt in a day or two. Always offer fresh milk replacer. Feeding equipment should be cleaned each day.

GENERAL RECOMMENDATIONS

1. Keep all equipment used for mixing and feeding thoroughly cleaned.
2. Newborn kids should receive colostrum as soon after birth as possible (within 3-4 hours).
3. When hand feeding by bottle or shallow pan, milk replacer should be about 100 to 105° F.
4. Consult with your veterinarian on a dry doe and kid vaccination program, including, but not limited to E. coli, clostridium and salmonella.
5. Always provide fresh, clean, free choice water and provide a high quality dry feed on a free choice basis.
6. If scouring occurs due to bacterial infection, kids may need treatment with an oral antibiotic. If scours persist, seek assistance from your local veterinarian.

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feeds.

Store in cool, dry location. Do not double stack pallets of milk replacer.

NOT FOR HUMAN CONSUMPTION



1495020-103

108

Manufactured By:

Land O'Lakes Animal Milk Products Co.
SHOREVIEW, MN 55126-2910

NET WT 25 LB (11.33 kg)