

Net Weight 33.3 Lbs. (15.12 Kgs.)

NO. 13413205

CATTLE BLOCK RU-MIN

(monensin Type C free-choice medicated feed)

Indications for Use

For increased rate of weight gain in growing beef steers and heifers on pasture (stocker, feeder, and slaughter) and replacement beef heifers.

FEED AS DIRECTED ON THIS LABEL

Active Drug Ingredient

Monensin 800 g/ton

Guaranteed Analysis

Crude Protein (Min.)	4.0	%
Crude Fat (Min.)	1.0	%
Crude Fiber (Max.)	12.0	%
Calcium (Min.)	4.3	%
Calcium (Max.)	5.3	%
Phosphorus (Min.)	2.0	%
Magnesium (Min.)	2.0	%
Salt (Min.)	11.0	%
Salt (Max.)	13.0	%
Copper (Min.)	200.0	ppm
Selenium (Min.)	12.0	ppm
Zinc (Min.)	800.0	ppm
Vitamin A (Min.)	120,000.0	IU/lb

Ingredients

plant protein products, roughage products, molasses products, calcium carbonate, dicalcium phosphate, salt, calcium hydroxide, potassium hydroxide, magnesium hydroxide, magnesium oxide, potassium sulfate, magnesium sulfate, manganous oxide, zinc oxide, ferrous sulfate, ferrous carbonate, copper sulfate, cobalt carbonate, sodium selenite, ethylenediamine, iron oxide, vitamin A supplement, feed grade fat product, sodium salts of fatty acids, hydrogenated tallow glyceride, precipitated chalk, sodium acid pyrophosphate, locust bean gum, karaya gum, anethole.

Feeding Directions

1. Feed this block free choice. Provide at least one block per 5 head of cattle.
2. Locate the blocks near the animal's feeding, watering or loafing areas.
3. Cattle should consume from 2 oz. to 8 oz. of block per head daily providing 50 mg to 200 mg of monensin per head daily. Discontinue feeding if block consumption falls below 2 oz. or rises above 8 oz daily.
4. Feed blocks continuously. Add one additional block when each block has been one-half consumed.
5. DO NOT feed salt or any other mineral supplement along with this block.

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.
 A withdrawal period has not been established for this product in pre-ruminating calves.
 Do not use in calves to be processed for veal.

Animal Safety Warnings

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

CAUTION: THIS FEED CONTAINS ADDED COPPER, DO NOT FEED TO SHEEP OR OTHER RELATED SPECIES.

Manufactured By:
Hi-Pro Feeds LLC
P.O. Box 519
Friona, Texas 79035



Approved by FDA under NADA # 119-253